

COVID-19 TRIAGE WORKSHEET

<p>SCREEN LOW RISK - See guidance below</p>	<p>ADMIT</p>	<p>TRANSFER</p>
<ul style="list-style-type: none"> <input type="checkbox"/> Screened or pending COVID-19 test <input type="checkbox"/> No history of exposure <input type="checkbox"/> Front line worker with URI symptoms/fever <input type="checkbox"/> Minimal risk factors as identified below <p>TREATMENT GUIDELINES</p> <ul style="list-style-type: none"> <input type="checkbox"/> Discharge home with supportive care recommendations and regular follow-up via phone. Return to ED if sx's worsen <p>Notify them of test results and recommend they remain self-quarantined until results back</p>	<ul style="list-style-type: none"> <input type="checkbox"/> Admit if concerning symptoms (SOA, dyspnea, wheeze, worsening condition Negative or PUI COVID-19 <input type="checkbox"/> Unable to maintain at home – ex: O2 sats with less than 5 l/min by NC or maintaining O2 sats >92% on non-rebreathing face mask. Minimal risk factors for SEVERE COVID-19 with cough/fever/myalgias. <p>(May show signs of sepsis with abnormal labs indicating organ dysfunction.)</p> <p>TREATMENT GUIDELINES</p> <ul style="list-style-type: none"> <input type="checkbox"/> Admit locally with sepsis protocol guidance & our COVID-19 order set. <p>Consider Hydroxychloroquine +/- azithromycin +/- zinc supplement</p> <p>Failure to improve or signs of worsening respiratory status – prepare for immed. transfer</p>	<ul style="list-style-type: none"> <input type="checkbox"/> COVID-19 positive or proven COVID-19 with High Risk Factors in respiratory distress. <p>*Respiratory rate is 35-40 breaths per minute, *Unable to maintain O2 sats >92% and/or PaO2 <65 on 6 l/min NC or NRB * Non-COPD with PaCO2 >50 with pH < 7.3</p> <ul style="list-style-type: none"> <input type="checkbox"/> Any patient requiring ventilatory support. <p>TREATMENT GUIDELINES</p> <p>See Surviving Sepsis Campaign guidance on “Summary of recommendations on the management of patients with COVID-19 and ARDS”</p>
<p>Patient-related risks</p> <p>Age > 60 -Immunocompromised (oncology, transplant, immunosuppressive meds, HIV, other known immunodeficiency) -Pregnancy - Chronic lung disease -Cirrhosis -Cardiovascular disease -End stage renal disease -Diabetes</p>	<p>Environmental risks</p> <p>-High-risk travel or known COVID-19 exposure within 14 days</p> <p>-Health care workers</p> <p>-Institutional home setting (nursing home, dormitory, shelter, prison, etc), outpatient dialysis center patient -Living with immunocompromised person(s)</p>	